



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

146. Proposed by L. C. WALKER, A.M., Professor of Mathematics, Colorado School of Mines, Golden, Col.

A random straight line crosses a given ellipse; find the chance that two points, taken at random in the ellipse, shall lie on opposite sides of the line.

Solution by G. B. M. ZEER, A. M., Ph. D., Parsons, W. Va.

Let  $O$  be the center of the given ellipse, axes  $OB=a$ ,  $OC=b$ . Let  $KL$  be the random chord,  $GH$  a diameter parallel to  $KL$ , and  $FE$  a diameter conjugate to  $GH$  meeting  $KL$  in  $Q$ . Let  $OG=n$ ,  $OE=m$ ,  $OQ=z$ ,  $\angle GOE=\phi$ ,  $\angle BOE=\theta$ . Equation to ellipse is  $x^2/m^2 + y^2/n^2 = 1$ .

Area of segment  $KEBL = A = 2\sin\phi \int y dx$ .

$$\therefore A = (2n\sin\phi/m) \int_z^m \sqrt{(m^2 - x^2)} dx = \frac{n\sin\phi}{2m} [\pi m^2 - 2z\sqrt{(m^2 - z^2)} - 2m^2 \sin^{-1}(z/m)]$$

Let  $p$  be the chance that both points are on the same side of  $KL$ ,  $p'$  the required chance. Also,  $\sin\phi = ab/mn$ .

$$\begin{aligned} \therefore p &= \int_0^m \int_{-m}^m 2A^2 dmdz / \pi^2 a^2 b^2 \int_0^m \int_{-m}^m dmdz \\ &= \frac{1}{4\pi^2 m^4} \int_0^m \int_{-m}^m \left[ \pi^2 m^4 + 4z^2(m^2 - z^2) + 4m^4 \left( \sin^{-1} \frac{z}{m} \right)^2 - 4\pi m^2 z \sqrt{(m^2 - z^2)} \right. \\ &\quad \left. - 4\pi m^4 \sin^{-1} \frac{z}{m} + 8m^2 z \sqrt{(m^2 - z^2)} \sin^{-1} \frac{z}{m} \right] dmdz \div \int_0^m m dm \\ &= \left( 1 - \frac{128}{45\pi^2} \right) \int_0^m m dm \div \int_0^m m dm = 1 - \frac{128}{45\pi^2}. \end{aligned}$$

$\therefore p' = 1 - p = 128/45\pi^2$ , the same as for a circle.

147. Proposed by W. J. GREENSTREET, A. M., Editor of The Mathematical Gazette, Stroud, England.

In a bag are  $n$  balls, known to be black or white, either color, *a priori*, equally likely. I draw two, which turn out to be one white and one black. I replace them and draw two more. What is the chance both are black?

Solution by W. W. LANDIS, Dickinson College, Carlisle, Pa.

This problem is open to two interpretations. We shall assume that a bag has been chosen at random from  $n+1$  bags containing respectively, 0, 1, 2, ...,  $n$  black balls. The chance of any particular number of black and white balls being found is then  $1/(n+1)$ . The chance that the bag containing  $r$  black balls is selected and a white and black ball drawn from it then  $1/(n+1) \cdot 2.r/n \cdot (n-r)/(n-1)$ . Adding these probabilities from  $r=1$  to  $r=n-1$  we obtain as sum  $\frac{1}{3}$ . As this event has happened, the probabilities in the next case must be multiplied by 3. We get then for the probability of drawing two black balls from this bag

$$\frac{6}{n+1} \cdot \frac{r}{n} \cdot \frac{n-r}{n-1} \cdot \frac{r}{n} \cdot \frac{r-1}{n-1} = 6[(n+1)r^3 - r^4 - nr^2]/(n+1)n^2(n-1)^2.$$